


1 **Postmortem Antigenic Epitope Retention of Galactose-N-Acetylglucosamine**
2 **in *Bacillus anthracis*-Infected Rhesus Monkey (*Macaca mulatta*) Spleens**
3
4
5
6
7

8 NEAL WOOLLEN¹, NANCY JAAX¹, WADE LAWRENCE¹, JOHN EZZELL², KELLY DAVIS¹,
9 AND ALFRED DENN¹



10
11
12
13
14
15 *Pathology Division¹ and Bacteriology Division²,*
16 *U.S. Army Medical Research Institute of Infectious Diseases,*
17 *Fort Detrick, Frederick, Maryland 21702-5011*
18

19950322 087

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

Galley proofs, correspondence, and requests for reprints should be sent to: Dr. Nancy Jaax, Pathology Division, U.S. Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, Maryland 21702-5011.

Running title: Immunohistochemical Staining of *Bacillus anthracis*

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract

BACKGROUND: Anthrax bacilli can be identified by using a monoclonal antibody to galactose-N-acetylglucosamine, a *Bacillus anthracis* cell-wall polysaccharide. We characterized the retention of antigenicity for this polysaccharide in prolonged postmortem autolysis of spleens from anthrax infected monkeys.

EXPERIMENTAL DESIGN: Spleens from infected rhesus monkeys were allowed to autolyze for up to 30 days. To visualize the availability and retention of the polysaccharide at various days of postmortem autolysis, splenic tissue impressions were immunostained by silver enhanced immunogold (IGSS) with a monoclonal antibody against galactose-N-acetylglucosamine.

RESULTS: All tissues had bacilli and fragments of bacilli present from the day of necropsy through 30 days of postmortem observation. Consistent immunocytochemical staining of bacilli occurred from days 3 to 15. In some cases, bacilli stained positively as late as 30 days. Extracellular stain reaction product demonstrating specific immunoreactivity was heavy in day 0 samples and gradually diminished over time to the day 15 samples.

CONCLUSIONS: The antigenic epitope recognized by the monoclonal antibody against galactose-N-acetylglucosamine was preserved with consistency for 15 days postmortem in decaying splenic tissue. Membrane-bound polysaccharide did not appear to be consistently available to antibody for reaction product formation until 3 days postmortem. The polysaccharide also appeared to exist free of the bacillus cell wall, and showed strong immunostaining properties during early postmortem decay.

Additional key words: Monoclonal Antibody, Galactose-N-acetylglucosamine, *Bacillus anthracis*

INTRODUCTION

Postmortem autolysis can hinder the definitive postmortem diagnosis of anthrax in humans, as well as in domestic and wild animals. In a publication describing 42 human cases of inhalation anthrax, anthrax bacilli were recovered by culture in only 20 cases. Twenty cases were culture negative and 2 were not examined. Histology on 15 of the culture-negative cases revealed bacilli (1). Theoretically, the availability of a reliable immunohistologic technique would have facilitated a definitive diagnosis in those 15 cases.

Antibodies have been produced to a variety of *Bacillus anthracis* antigenic subunits. Antibody to a *B. anthracis* cell-wall polysaccharide, galactose-N-acetylglucosamine, has demonstrated a high level of specificity, suggesting it has immunodiagnostic potential (2). It demonstrated superior immunohistochemical signal producing potential in a preliminary screen that also included antibodies to a toxin protein, capsular peptide (poly-D-glutamic acid), and cell-wall protein. However, preservation of the polysaccharide's antigenicity during autolysis has not been reported. In this paper, we report the distribution, availability to antibody, and preservation of the antigenic galactose-N-acetylglucosamine polysaccharide during autolysis of infected tissue.

EXPERIMENTAL DESIGN

Spleen samples from 5 rhesus monkeys infected with the Vollum 1B strain of *B. anthracis*, which had previously served as controls in a drug and vaccine efficacy study (3), were allowed to autolyze for up to 30 days at room temperature (20 to 22° C). Tissue impressions made at days 0, 1, 2, 3, 4, 5, 6, 9, 11, 13, 15, 18, 21, and 30 were Giemsa stained (Baxter, Deerfield, IL) and immunostained by silver enhanced immunogold (IGSS) with a mouse monoclonal antibody against the cell-wall polysaccharide, galactose-N-acetylglucosamine. Pattern and intensity of positive immunostain reaction product were analyzed for all samples.

RESULTS AND DISCUSSION

Two distribution patterns of the IGSS reaction product were observed. The reaction product was either closely bacillus-associated (b/a) or free in the extracellular tissue fluid (e/c) showing no

1 direct association with bacilli. Mean values for both Giemsa- and IGSS-stained impressions are
2 displayed in Figure 1.

3 The b/a signal, initially negligible, was strong by 3 days of tissue autolysis. This suggests that
4 the organism-bound polysaccharide was not accessible to antibody binding and/or reaction product
5 formation in fresh tissue, but became increasingly accessible with postmortem autolysis, possibly
6 due to the destruction of the organism's protective poly-D-glutamic acid capsule. By day 3, there
7 were 1 to 20 IGSS-stained organisms per high-power field (org/hpf) on impressions of tissues
8 from all animals. Consistent staining of bacilli was noted from days 3 to 15. Before day 3, IGSS-
9 stained bacilli had an irregular stain pattern (Fig. 2). From day 3 to 9, cell walls of the IGSS-
10 stained bacilli had a uniform stain pattern (Fig. 3), which suggested that antigen was easily
11 accessible to the primary antibody for reaction product formation. Again we noted an irregular cell
12 wall stain pattern in IGSS-stained organisms from tissues subjected to 11 or more days of
13 autolysis. From days 3 to 21, there were equal to or greater numbers of bacilli observed on IGSS-
14 stained impressions than on Giemsa-stained impressions, which suggested that immunostaining
15 with anti-galactose-N-acetylglucosamine enhanced recognition of anthrax bacilli over this period of
16 tissue autolysis. The number of Giemsa-stained microorganisms, < 1 org/hpf on day 18
17 impressions, gradually rose on days 21 and 30, surpassing the number of IGSS-stained
18 microorganisms on the day 30 impressions. This most likely represents proliferation of
19 contaminating bacterial growth, as significant additional sporulation and new growth of anthrax
20 bacilli would not be anticipated in the airtight microenvironment. In addition to specificity testing
21 described in prior work (2), we tested for and found no cross-reactivity of the primary antibody
22 with *Clostridium sporogenes*, *B. subtilis*, *Escherichia coli*, *Salmonella enteritidis*, *Pasteurella*
23 *multocida*, *Yersinia enterocolitica*, *Bordetella bronchiseptica*, *Brucella abortus*, *Brucella melitensis*,
24 and *Brucella suis*.

25 The amount of e/c reaction product, initially heavy, lessened with prolonged tissue autolysis.
26 The heavy concentration on day 0 impressions persisted with no remarkable change until day 3.

1 From day 3 to 15, reduced yet notable e/c reaction product persisted. With rare exception, there
2 was no remarkable e/c signal from day 18 to 30. This pattern of staining was never observed in
3 stain controls testing for nonspecific binding of the secondary antibody or on impressions prepared
4 from non-infected monkey spleens. This suggests that it was a positive indication of non-cellular
5 bound galactose-N-acetylglucosamine free in *B. anthracis*-infected tissue and fluids.

6 METHODS

7 SAMPLE COLLECTION AND SLIDE PREPARATION

8 Spleen samples for day 0 tissue impressions were taken from 5 untreated *B. anthracis*-infected
9 rhesus monkeys (*Macaca mulatta*) immediately postmortem (3). After gross necropsy
10 observations and splenic sampling, body cavities were closed and the carcasses were allowed to
11 remain at room temperature (20 to 22° C) for 24 hr. Using aseptic techniques, we harvested
12 spleens, diced them into approximate 0.5 cm cubes, and placed them in small capped containers to
13 allow room temperature autolysis for 2, 3, 4, 5, 6, 9, 11, 13, 15, 18, 21, and 30 days. Day 1
14 tissue impressions made at this time were placed on Superfrost/Plus (Fisher Scientific, St. Louis,
15 MO) microscope slides, air dried, and fixed for 1 hr in 10% buffered neutral formalin. One slide
16 from each group was Giemsa stained to evaluate the relative number of bacilli present.

17 IMMUNOSTAIN TECHNIQUE

18 After formalin fixation, the impressions were processed by the following method: a) reverse
19 osmosis (r/o) water, 2 x 5 min; b) Lugol's iodine (4), 1 x 5 min; c) r/o water, 1 x 5 min; d) 5%
20 sodium thiosulfate, 15 sec rinse (4); e) r/o water, 3 x 5 min; f) wash buffer consisting of 0.8%
21 bovine serum albumin (BSA) (Sigma Chemical Company, St. Louis, MO), 0.1% fish skin gelatin
22 (FSG) (Sigma Chemical Company), 0.5% triton-X 100 (Sigma Chemical Company), 0.02 M
23 glycine (Sigma Chemical Company), and 2 mM sodium azide (Sigma Chemical Company) in 0.05
24 M tris-buffered saline (TBS), 2 x 5 min (5); and g) blocking buffer consisting of 0.08% BSA,
25 0.01% FSG, 5% goat serum (Sigma Chemical Company), and 2 mM sodium azide in 0.05 M
26 TBS, 1 x 15 min. Impressions were drained without rinsing and treated with MAb EAII-6G6 (2)

1 primary antibody diluted 1:100 in incubating buffer consisting of 0.08% BSA, 0.1% FSG, 1%
2 goat serum, 0.1% triton-X 100, and 2 mM sodium azide in 0.05 M TBS for 90 min; washed with
3 buffer, 2 x 15 min; and then treated with blocking buffer, 1 x 10 min. Impressions were again
4 drained without rinsing and 5 nm gold conjugated goat anti-mouse secondary antibody (Amersham
5 Corporation, Arlington Heights, IL) diluted 1:40 in incubating buffer was applied for 60 min;
6 followed by wash buffer, 3 x 5 min; r/o water, 30 sec rinse; r/o water, 4 x 5 min; and silver
7 enhancement (Amersham Corporation), approximately 10 min.

8 If the silver autprecipitation time was < 20 min, the silver enhancement step was split into two
9 5 min treatments. The impressions were counterstained with working light green solution (6).

10 Nonspecific binding of the secondary antibody was tested for by eliminating the primary antibody
11 application and subsequent steps, proceeding immediately to application of the secondary
12 antibody. Nonspecific binding of the primary antibody was tested for by staining normal monkey
13 spleen impressions at 0, 5, 7, and 9 days of autolysis. A known anthrax positive spleen
14 impression control was also processed with this group. When a day group stained either negative
15 or variable, the group was restained along with a positive control to confirm that the results were
16 not due to a procedural error.

17 SLIDE INTERPRETATION

18 Stained impressions were evaluated by brightfield microscopy. The formation of a
19 brown-black reaction product indicated the presence of the target antigen. A subjective score of 0
20 (negligible) to 4 (heavy) was assigned for each impression, characterizing the amount of e/c
21 reaction product. Scores of 0 to 4 were also assigned for evaluating the number of Giemsa-stained
22 bacilli as well as evaluation of the b/a signal based on 0 < 1 org/hpf, 1 = 1-20 org/hpf, 2 = 21-60
23 org/hpf, 3 = 61-100 org/hpf, and 4 > 100 org/hpf. The final score assigned for each day of
24 sample evaluation was based on the average for the 5 animals. High-power field (hpf) is defined,
25 in this study, as microscopy with a 40X objective and 10X eyepieces (400X).

26 TEST FOR BACTERIAL CROSS-REACTIVITY

1 Using the same immunostain technique described above, we stained formalin-fixed spot slides
2 of *C. sporogenes*, and used formalin-fixed *B. anthracis* (Sterne strain) spot slides as positive
3 controls.

4 Using the monoclonal antibody labeled with fluorescein isothiocyanate (FITC), we tested for
5 additional cross-reactivity using formalin-fixed spot slides of the following bacteria: *B. subtilis*,
6 *C. sporogenes*, *E. coli*, *S. enteritidis*, *P. multocida*, *Y. enterocolitica*, *B. bronchiseptica*, *Br.*
7 *abortus*, *Br. melitensis*, and *Br. suis*. Impressions of day 3 postmortem *B. anthracis*-infected
8 spleens served as positive controls. Briefly, the impressions were rinsed in distilled water,
9 incubated at room temperature in a humidified chamber for 30 min with normal mouse serum
10 diluted 1:200 in phosphate-buffered saline (PBS), pH 7.4, followed by a 30-min incubation of
11 FITC-labeled monoclonal antibody to galactose-N-acetylglucosamine diluted 1:5 in PBS. The
12 impressions were rinsed in PBS, then in distilled water, mounted with Vectashield mounting
13 medium (Vector Laboratories, Burlingame, CA), and examined by fluorescence microscopy.
14

REFERENCES

1. Abramova FA, Grinberg LM, Yampolskaya OV, Walker DH. Pathology of inhalational anthrax in 42 cases from the Sverdlovsk outbreak of 1979. *Proc Natl Acad Sci* 1993;90:2291-4.
2. Ezzell Jr JW, Abshire TG, Little SF, Lidgerding BC, Brown C. Identification of *Bacillus anthracis* by using monoclonal antibody to cell wall galactose-*N*-acetylglucosamine polysaccharide. *J Clin Microbiol* 1990;28(2):223-31.
3. Friedlander AM, Welkos SL, Pitt MLM, Ezzell JW, Worsham PL, Rose KJ, et al. Postexposure prophylaxis against experimental inhalation anthrax. *J Infect Dis* 1993;167:1239-42.
4. Immunogold staining with silver enhancement (IGSS). In: Filipe MI, Lake BD, editors. *Histochemistry in pathology*. New York; Churchill Livingstone, 1990:480-1.
5. Larsson LI. Buffers and detergents. In: *Immunocytochemistry: Theory and practice*. Boca Raton, Florida; CRC Press, Inc., 1988:160-1.
6. Luna LG. Methods for carbohydrates and mucoproteins. In: *Manual of histologic staining methods of the Armed Forces Institute of Pathology*. New York; McGraw-Hill Book Company, 1968:159.

1 FIG. 1. A comparison of mean score values for the relative numbers of Giemsa- and IGSS-
2 stained bacilli (b/a) and amount of extracellular (e/c) IGSS stain reaction product. A subjective
3 score of 0 (negligible) to 4 (heavy) was assigned for each impression. Scores of 0 to 4 were also
4 assigned for evaluating the number of Giemsa-stained bacilli as well as evaluation of the b/a signal
5 based on 0 < 1 org/hpf, 1 = 1-20 org/hpf, 2 = 21-60 org/hpf, 3 = 61-100 org/hpf, and 4 > 100
6 org/hpf.

7 FIG. 2. An IGSS-stained chain of bacilli from a spleen with 2 days of postmortem autolysis.
8 The stain pattern for bacilli was irregular and extracellular stain reaction product was moderately
9 heavy (score=3), X1,800.

10 FIG. 3. An IGSS-stained chain of bacilli from a spleen with 3 days of postmortem autolysis.
11 The stain pattern for bacilli was uniform and the average extracellular stain reaction product was
12 moderate (score=2). In this micrograph, extracellular stain reaction product is mild (score=1),
13 X1,400.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution unlimited - approved for public release		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION US Army Medical Research Institute of Infectious Disease		6b. OFFICE SYMBOL (If applicable) MCMR-UIP-C	7a. NAME OF MONITORING ORGANIZATION US Army Medical Research and Development Command		
6c. ADDRESS (City, State, and ZIP Code) Ft. Detrick Frederick, MD 21702-5011			7b. ADDRESS (City, State, and ZIP Code) Ft. Detrick Frederick, MD 21702-5012		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
11. TITLE (Include Security Classification) Postmortem Antigenic Epitope Retention of Galactose-N-Aceylglucosamine in Bacillus anthracis-Infected Rhesus Monkey (Macaca mulatta) Spleens					
12. PERSONAL AUTHOR(S) Neal Woollen, Nancy Jaax, Wade Lawrence, John Ezzell, Kelly Davis, Alfred Denn					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 23 Nov 94	
15. PAGE COUNT 11					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) BACKGROUND: Anthrax bacilli can be identified by using a monoclonal antibody to galactose-N-acetylglucosamine, a Bacillus anthracis cell wall polysaccharide. We characterized the retention of antigenicity for this polysaccharide in prolonged post-mortem autolysis of spleen from infected monkeys. EXPERIMENTAL DESIGN: Spleens from infected rhesus monkeys were allowed to autolyze for up to 30 days. To visualize the availability and retention of the polysaccharide at various days of postmortem autolysis, splenic tissue impressions were immunostained by silver enhanced immunogold (IGSS) using a monoclonal antibody against galactose-N-acetylglucosamine. RESULTS: All tissue had bacilli & fragments of bacilli present from the day of necropsy through 30 days postmortem observation. Consistent immunocytochemical staining of bacilli occurred from day 3 to 15. In some cases, bacilli stained positive as late as 30 days. Extracellular stain reaction product demonstrating specific immunoreactivity was also noted as heavy in day 0 samples and gradually diminished over time to the day 15 samples. CONCLUSIONS: The antigenic epitope recognized by the monoclonal antibody against galactose-N-acetylglucosamine was preserved with consistency for 15 days postmortem in decaying splenic tissue. Membrane bound polysaccharide did not appear to be consistently available to antibody and reaction product formation until 3 days postmortem. The polysaccharide appears to also exit free of the bacillus cell wall, & showed strong immunostaining properties during early post-mortem decay.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL